

A LIST OF BIRDS TAKEN ON MARAGUANA,
WATLING'S ISLAND, AND INAGUA. BAHIA-
MAS, DURING JULY, AUGUST, SEPTEMBER,
AND OCTOBER, 1891.

BY CHARLES B. CORY.

MARAGUANA (August 5 to September 14).

| | |
|--|--|
| <i>Larus atricilla</i> Linn. | <i>Ceryle alcyon</i> (Linn.). |
| <i>Gelochelidon nilotica</i> (Hasselq.). | <i>Tyrannus dominicensis</i> (Gmel.). |
| <i>Ardea tricolor ruficollis</i> Gosse. | <i>Loxigilla violacea</i> (Linn.). |
| <i>Ardea virescens</i> (Linn.). | <i>Spindalis zena</i> (Linn.). |
| <i>Nycticorax violaceus</i> (Linn.). | <i>Euetheia bicolor</i> (Linn.). |
| <i>Himantopus mexicanus</i> (Müll.). | <i>Chelidon erythrogaster</i> (Bodd.). |
| <i>Tringa maculata</i> Vieill. | <i>Vireo crassirostris</i> (Bryant). |
| <i>Tringa minutilla</i> Vieill. | <i>Cœreba bahamensis</i> (Reich.). |
| <i>Totanus flavipes</i> (Gmel.). | <i>Dendroica petechia gundlachi</i> |
| <i>Symphemia semipalmata</i> (Gmel.). | <i>Baird.</i> |
| <i>Bartramia longicauda</i> (Bechst.). | <i>Dendroica dominica</i> (Linn.). |
| <i>Ægialitis vocifera</i> (Linn.). | <i>Dendroica discolor</i> (Vieill.). |
| <i>Ægialitis wilsonia</i> (Ord). | <i>Mimus polyglottus orpheus</i> |
| <i>Arenaria interpres</i> (Linn.). | <i>(Linn.).</i> |
| <i>Hæmatopus palliatus</i> Temm. | <i>Mimus gundlachi</i> Caban. |
| <i>Columba leucocephala</i> (Linn.). | <i>Polioptila cærulea</i> (Linn.). |
| <i>Columbigallina passerina</i> (Linn.). | <i>Margarops fuscatus</i> (Vieill.). |

WATLING'S ISLAND (September 28 to November 9).

| |
|---|
| <i>Larus atricilla</i> (Linn.). |
| <i>Phalacrocorax dilophus floridanus</i> (Aud.). — Eight specimens. |
| <i>Ardea tricolor ruficollis</i> (Gosse). |
| <i>Ægialitis vocifera</i> (Linn.). |
| <i>Coccyzus minor maynardi</i> Ridgw. |
| <i>Coccyzus americanus</i> (Linn.). |
| <i>Tyrannus dominicensis</i> (Gmel.). |
| <i>Dolichonyx oryzivorus</i> (Linn.). — Oct. 12, female. |
| <i>Habia ludoviciana</i> (Linn.). — Oct. 20, female. |
| <i>Euetheia bicolor</i> (Linn.). |
| <i>Vireo olivaceus</i> (Linn.). — Oct. 5, male. |
| <i>Cœreba bahamensis</i> (Reich.). |
| <i>Mniotilta varia</i> (Linn.). |
| <i>Compsothlypis americana</i> (Linn.). — Five specimens. |

- Dendroica tigrina* (Gmel.). — Seven specimens.
Dendroica petechia gundlachi Baird. — Common, breeds.
Dendroica maculosa (Gmel.). — Six specimens, Oct. 6-21.
Dendroica striata (Forst.). — Thirty-six specimens.
Dendroica blackburniæ (Gmel.). — Two specimens, Oct. 12.
Dendroica dominica (Linn.). — Six specimens.
Dendroica virens (Gmel.). — Oct. 12, male.
Dendroica palmarum (Gmel.). — Oct. 20, female.
Dendroica discolor (Viell.). — Twenty-five specimens.
Seiurus aurocapillus (Linn.). — Thirteen specimens.
Mimus gundlachi Caban.
Margarops fuscatus (Viell.).

INAGUA.

- Gallinula galeata* (Licht.). — July 17.
Tringa minutilla Viell. — July 28.
Totanus flavipes (Gmel.). — July 28.
Myiarchus sagræ Gundl.
Loxigilla violacea (Linn.).
Vireo olivaceus (Linn.). — Sept. 17, female.
Vireo crassirostris (Bryant).
Cœreba bahamensis (Reich.).
Helmitherus vermivorus (Gmel.). — Sept. 22, female.
Dendroica petechia gundlachi Baird.
Mimus gundlachi Caban.

 NOTES ON THE RANGE AND HABITS OF THE
 CAROLINA PARRAKEET.

BY AMOS W. BUTLER.

THE CAROLINA PARRAKEET (*Conurus carolinensis*), whose range is now confined to quite restricted areas in some of our southern States, was formerly known as a characteristic bird of Indiana. At the time of its greatest range in that State, within historic times, it was known from New York, Pennsylvania and Maryland to Kansas, Nebraska, and possibly Colorado. It is my desire to present some evidence tending to show its distribution in Indiana and neighboring States together with some notes upon its habits.